



Figure 1

**Examples of Substituted Piperidines Accessible Via
the Methods of the Present Invention**



Structure A

Structure B



Structure C

Structure D

For Structures A, B, C, and D:

$n = 0, 1, \text{ or } 2$

$R_1 = \text{H, alkyl, aryl, heteroaryl, aralkyl, } -\text{CO}_2R_4, \text{ or } -\text{C(O)NHR}_4$

$R_2 = \text{alkyl}$

$X = \text{O, NC(O)R}_4, \text{ or NS(O)}_2R_4, \text{ NH, NR}_4, \text{ S, or S(O)}$

$R_3 = \text{alkyl, aralkyl, aryl, or heteroaryl}$


$R_4 = \text{alkyl, aryl, heteroaryl, or aralkyl}$

(



Figure 3

Prophetic Asymmetric Synthesis of **50**



50

Figure 4

Prophetic Asymmetric Syntheses of 3-Substituted Piperidine 1






Figure 5

Prophetic Asymmetric Synthesis of 3-Substituted Piperidine 3




TADDOL catalysts




See Seebach, D. et al.
Tetrahedron **1992**, 48 (27), 5719.

Figure 6






Prophetic Asymmetric Synthesis of 3-Substituted Piperidine **51**



TADDOL catalysts



See Seebach, D. et al.
Tetrahedron **1992**, 48 (27), 5719.






Figure 9




Figure 10




Figure 11

Mixture of 15, 16, 27, and 28




Figure 12

15: (R,S)-Isomer Chromatogram




Figure 13

27: *(S,S)*-Isomer Chromatogram




Figure 14

16: (R,R)-Isomer Chromatogram




Figure 15

28: (S,R)-Isomer Chromatogram




Figure 16

No.	Description of HPLC Trace	Peak Retention Times (min)
1	Catalyst 13 (254 nm)	8.420, 8.781
2	Catalyst 13 (220 nm)	8.420, 8.781
3	Catalyst 13 and non-selective coinjection (254 nm)	8.155, 8.328, 8.695
4	Catalyst 13 and non-selective coinjection (220 nm)	8.155, 8.333, 8.688
5	non-selective (254 nm)	8.208, 8.395, 8.688
6	Catalyst 14 (254 nm)	8.061, 8.210, 8.399, 8.688, 8.897

Figure 17

No.	Description of HPLC Trace	Peak Retention Times (min)
1	Catalyst 14 (254 nm)	8.158, 8.423
2	Catalyst 14 (220 nm)	8.030, 8.159, 8.366
3	Catalyst 14 and non-selective coinjection (254 nm)	8.176, 8.386, 8.664
4	Catalyst 14 and non-selective coinjection (220 nm)	8.044, 8.178, 8.387
5	non-selective (254 nm)	8.176, 8.374, 8.646, 9.950
6	non-selective (220 nm)	8.375

Figure 18

No.	Description of HPLC Trace	Peak Retention Times (min)
1	Catalyst 13 (254 nm)	8.42
2	Catalyst 13 (220 nm)	8.420, 8.781
3	Catalyst 13 and Catalyst 14 coinjection (254 nm)	8.147, 8.337, 8.695
4	Catalyst 13 and Catalyst 14 coinjection (220 nm)	8.147, 8.338, 8.695
5	Catalyst 14 (254 nm)	8.158, 8.423
6	Catalyst 14 (220 nm)	8.030, 8.159, 8.366

Figure 19




Figure 20











Figure 21




Figure 22



















Figure 23




R = Me, TBDMS, TBDPS



R = Me, H, $-\text{CH}_2\text{CMe}_3$



R = Me, Et, Ph



R = H, $-(\text{CH}_2)_2\text{-2-pyridyl}$,
 $-(\text{CH}_2)\text{C(Ph)}_2\text{OH}$

Figure 24













Figure 25












Figure 26






Figure 27
















Figure 28

Figure 29

Aminoalcohols



R = n Bu, $-(CH_2)_4^-$,
 $-(CH_2)_5^-$, $-(CH_2)_6^-$






Figure 31

Figure 32











Figure 33










Figure 34












Figure 35



● = an organic polymer or inorganic solid support

Figure 36



() ()




















Figure 38




Figure 39




Figure 40

Mixture of 1, 2, 3 and 4:




Figure 41

2: (R,S)-Isomer Chromatogram

